Fractal Scales in a Schwarzschild Atmosphere

E.N. Glass* and J.P. Krisch

Department of Physics, University of Michigan, Ann Arbor, Michgan 48109

(18 May 2000)

arXiv:gr-qc/0006020v1 6 Jun 2000

Recently, Glass and Krisch have extended the Vaidya radiating metric to include both a radiation fluid and a string fluid [1999 Class. Quantum Grav. **16**, 1175]. Mass diffusion in the extended Schwarzschild atmosphere was studied. The continuous solutions of classical diffusive transport are believed to describe the envelope of underlying fractal behavior. In this work we examine the classical picture at scales on which fractal behavior might be evident.

PACS numbers: 04.20.Jb, 5.45Df, 64.6

*Permanent address: Physics Department, University of Windsor, Ontario N9B 3P4, Canada

I. INTRODUCTION

Since its discovery in 1916, the Schwarzschild vacuum solution has been a source of many insights into relativistic physics.

$$g_{ab}^{Sch}dx^a dx^b = (1 - 2m_0/r)du^2 + 2dudr - r^2 d\Omega^2. \tag{1}$$

This solution and its extensions contain new views of relativistic physics on a variety of scales, from the size of the Universe down to the size of a stellar black hole. When contemplating quantum effects at classical boundaries such as the Schwarzschild horizon, the scales become much smaller down to the Planck length $L_p = \sqrt{\hbar G/c^3} = 1.6 \times 10^{-33}\ cm$. The classical point particle electron may perhaps be resolved into Planck scale strings bits with size proportional to the Planck length and with 10^{20} strings bits fitting into a classical electron radius.

Vaidya [1] extended the solution by allowing the Schwarzschild mass parameter to be a function of retarded time, $m(u)$, and discovered a spherically symmetric null fluid atmosphere beyond the horizon at $2m_0$. Glass and Krisch (GK) [2] pointed out that allowing the Schwarzschild mass parameter to be a function of radial position and retarded time, $m(u, r)$, creates an atmosphere with both a null fluid and string fluid stress-energy content described by metric g_{ab}^{GK}. The density of the string fluid, ρ, obeys a classical diffusion equation in the Schwarzschild spacetime:

$$\partial_u \rho = r^{-2}\partial_r(Dr^2\partial_r\rho) \tag{2}$$

where $D(r)$ is the variable diffusivity [3] found in anomalous diffusion. One may show [4], if the mass is diffusing in a flowspace with $\sqrt{-g} = r^{-\beta}f(\vartheta)$ (in [4] $\beta = 1 - \delta$), that the functional dependence of the diffusivity is determined.

$$\partial_u m = r^{\beta}\partial_r(Dr^{-\beta}\partial_r m) \tag{3}$$

$$D(r) = D_0 r^{\beta-2}.$$

The continuous solutions of classical diffusive transport are believed to describe the envelope of underlying fractal behavior [5]. Fractal behavior is usually associated with mass scaling but the radial dependence of the diffusivity [6] carries additional information about the connectivity of the underlying fractal substrate, and so we wish to examine the classical picture at scales on which fractal behavior might be evident. These smaller scales are still classical (i.e. larger than quantum scales). We consider a 2-parameter family of spacetimes with metric function $A = 1 - 2m(u,r)/r$

$$g_{ab}^{scale} dx^a dx^b = A du^2 + 2 du dr - \lambda^{2-\alpha} r^\alpha d\Omega^2, \tag{4}$$

where λ has units of length, and is required for spheres to have area of (length)^2. This metric is capable of describing different scales for a particular central source. For example, a particular mass function $m(u,r)$ with $\alpha = 2$, could describe the Schwarzschild vacuum outside the source. For $\alpha \neq 2$ it would describe the source at different scales where we expect evidence of fractal behavior to appear. Fractal dimensions are generally less than the Euclidean dimension, and we will describe an averaging process on 2-spheres for $\alpha \leq 2$. The scale parameter α cannot be removed by a coordinate transformation. If one attempts the appearance of the usual area coordinate by $y^2 = \lambda^{2-\alpha} r^\alpha$, then α will emerge in the (u,r) part of g_{ab}^{scale}. Moreover, the manifestly coordinate-independent sectional curvature of the (ϑ, φ) two-surfaces, given in Eq.(15) below, provides a physically meaningful mass expression in terms of α.

The interpretation of α as a fractal scale follows from considering diffusive processes in the atmosphere. If $\mathcal{P}(u,r)$ is a quantity that is diffusing, the diffusion equation is

$$\partial_u \mathcal{P} = r^{-\alpha} \partial_r (D r^\alpha \partial_r \mathcal{P}) \tag{5}$$

and $\alpha + 1$ is identified as the dimension of the fractal substrate on which the diffusion is occurring [5], [7]. At smaller quantum levels, $\mathcal{P}$ can be a probability measure on the space of radial paths [8].

The fluid atmosphere described by this metric is discussed in the next section. In section

III diffusion and scale sizes are discussed. Definitions of topological and Hausdorff dimensions are given. Applications of the fractal view are presented in section IV, along with specific mass solutions. The paper is summarized in the final section.

In this work Latin indices range over (0,1,2,3) $= (u, r, \vartheta, \varphi)$. Our sign conventions are $2A_{c;[ab]} = A_e R^e{}_{cab}$, and $R_{ab} = R^c{}_{abc}$. Overdots abbreviate $\partial/\partial u$, and primes abbreviate $\partial/\partial r$. Overhead carets denote unit vectors. We use units where $G = c = 1$. Einstein's field equations are $G_{ab} = -8\pi T_{ab}$, and the metric signature is (+,-,-,-).

II. THE ATMOSPHERE

Metric g_{ab}^{scale} describes an atmosphere that admits a two-fluid description of matter with outward flowing short-wavelength photons (sometimes called a "null fluid").

With the use of a Newman-Penrose null tetrad the Einstein tensor for the atmosphere is computed from (4) and given by

$$G_{ab} = -2\Phi_{11}(l_a n_b + n_a l_b + m_a \bar{m}_b + \bar{m}_a m_b) \tag{6}$$
$$-2\Phi_{00} n_a n_b - 2\Phi_{22} l_a l_b - 6\Lambda g_{ab}.$$

Here the null tetrad components of the Ricci tensor are

$$\Phi_{00} = \frac{\alpha(2-\alpha)}{4r^2}, \tag{7a}$$
$$\Phi_{11} = \frac{2m' - rm''}{4r^2} - \frac{\alpha^2}{16r^2} + \frac{\lambda^{\alpha-2}}{4r^\alpha} + \frac{(\alpha-2)(\alpha+2)m}{8r^3}, \tag{7b}$$
$$\Phi_{22} = -\frac{\alpha\dot{m}}{2r^2} + \frac{\alpha(2-\alpha)}{16r^2}(1-2m/r)^2, \tag{7c}$$
$$\Lambda = R/24 = \frac{rm'' + 2(\alpha-1)m'}{12r^2} + \frac{(\alpha-2)(3\alpha-2)m}{24r^3} \tag{7d}$$
$$-\frac{\alpha(3\alpha-4)}{48r^2} + \frac{\lambda^{\alpha-2}}{12r^\alpha}.$$

The only non-zero component of the Weyl tensor is

$$\Psi_2 = -\frac{(\alpha+1)m}{3r^3} + \frac{(2+\alpha)m' - rm''}{6r^2} + \frac{\alpha}{12r^2} - \frac{\lambda^{\alpha-2}}{6r^\alpha}. \tag{8}$$

The metric is Petrov type **D** with l^a and n^a principal null geodesic vectors

$$l_a dx^a = du, \qquad l^a \partial_a = \partial_r \tag{9a}$$

$$n_a dx^a = (A/2)du + dr, \qquad n^a \partial_a = \partial_u - (A/2)\partial_r \tag{9b}$$

$$m_a dx^a = -(1/\sqrt{2})\lambda^{1-\alpha/2} r^{\alpha/2}(d\vartheta + i\,\sin\vartheta d\varphi), \quad m^a \partial_a = (1/\sqrt{2})\lambda^{\alpha/2-1} r^{-\alpha/2}(\partial_\vartheta + \frac{i}{\sin\vartheta}\partial_\varphi). \tag{9c}$$

When $\alpha = 2$ all quantities have the spacetime values for metric g_{ab}^{GK} discussed in [2].

In order to clearly see the matter content we introduce a time-like unit velocity vector $\hat{v}^a$ and three unit space-like vectors $\hat{r}^a$, $\hat{\vartheta}^a$, $\hat{\varphi}^a$ such that

$$g_{ab}^{scale} = \hat{v}_a \hat{v}_b - \hat{r}_a \hat{r}_b - \hat{\vartheta}_a \hat{\vartheta}_b - \hat{\varphi}_a \hat{\varphi}_b.$$

The unit vectors are defined by

$$\hat{v}_a dx^a = A^{1/2} du + A^{-1/2} dr, \qquad \hat{v}^a \partial_a = A^{-1/2}\partial_u, \tag{10a}$$

$$\hat{r}_a dxa = A^{-1/2} dr, \qquad \hat{r}^a \partial_a = A^{-1/2}\partial_u - A^{1/2}\partial_r, \tag{10b}$$

$$\hat{\vartheta}_a dx^a = \lambda^{1-\alpha/2} r^{\alpha/2} d\vartheta, \qquad \hat{\vartheta}^a \partial_a = -\lambda^{\alpha/2-1} r^{-\alpha/2}\partial_\vartheta, \tag{10c}$$

$$\hat{\varphi}_a dx^a = \lambda^{1-\alpha/2} r^{\alpha/2} \sin\vartheta d\varphi, \qquad \hat{\varphi}^a \partial_a = -\lambda^{\alpha/2-1}(r^{\alpha/2}\sin\vartheta)^{-1}\partial_\varphi. \tag{10d}$$

The Einstein tensor (6) yields the energy-momentum tensor

$$T_{ab} = \psi l_a l_b + \rho \hat{v}_a \hat{v}_b + p_r \hat{r}_a \hat{r}_b + p_\perp(\hat{\vartheta}_a \hat{\vartheta}_b + \hat{\varphi}_a \hat{\varphi}_b) + q_a \hat{v}_b + \hat{v}_a q_b \tag{11}$$

where

$$8\pi\psi = -\frac{\alpha \dot{m}}{r^2} + \frac{\alpha(2-\alpha)A^2}{8r^2}, \tag{12a}$$

$$8\pi\rho = \frac{\alpha m'}{r^2} + \frac{\lambda^{\alpha-2}}{r^\alpha} - \frac{\alpha(\alpha-1)}{2r^2} + \frac{(\alpha-2)}{8r^3}[8\alpha m - \alpha(r-2m)], \tag{12b}$$

$$8\pi p_r = -\frac{\alpha m'}{r^2} - \frac{\lambda^{\alpha-2}}{r^\alpha} + \frac{\alpha(\alpha-1)}{2r^2} - \frac{(\alpha-2)}{8r^3}[8\alpha m + \alpha(r-2m)], \tag{12c}$$

$$8\pi p_\perp = -\frac{m''}{r} - \frac{(\alpha-2)m'}{r^2} - \frac{(\alpha-2)^2 m}{2r^3} + \frac{\alpha(\alpha-2)}{4r^2}, \tag{12d}$$

$$8\pi q_a = [A\frac{\alpha(2-\alpha)}{8r^2}]\,\hat{r}_a. \tag{12e}$$

The string equation of state $\rho + p_r = 0$ holds for the scale with $\alpha = 2$. The decomposition of the energy-momentum tensor given in Eq.(11) is less compelling on $\alpha \neq 2$ scales since there

is heat flow in addition to the null fluid. The heat flow arises from the $\Phi_{00} n_a n_b$ term in G_{ab}. This is seen only on $\alpha \neq 2$ scales and is the intermediate electromagnetic field part of the null fluid, which has only a far field part in the usual Vaidya large scale view. A different covariant decomposition using $l_a dx^a = A^{-1}(\hat{v}_a - \hat{r}_a)dx^a$ provides a single fluid with radial heat flow. The energy-momentum tensor decomposes as

$$T_{ab} = \tilde{\rho}\hat{v}_a\hat{v}_b + \tilde{p}_r\hat{r}_a\hat{r}_b + p_\perp(\hat{\vartheta}_a\hat{\vartheta}_b + \hat{\varphi}_a\hat{\varphi}_b) + \tilde{q}_a\hat{v}_b + \hat{v}_a\tilde{q}_b \tag{13}$$

where

$$8\pi\tilde{\rho} = -A^{-1}\frac{\alpha\dot{m}}{r^2} + \frac{\alpha m'}{r^2} + \frac{\lambda^{\alpha-2}}{r^\alpha} - \frac{\alpha(\alpha-1)}{2r^2} + \frac{(\alpha-2)}{4r^3}[4\alpha m - \alpha(r-2m)], \tag{14a}$$

$$8\pi\tilde{p}_r = -A^{-1}\frac{\alpha\dot{m}}{r^2} - \frac{\alpha m'}{r^2} - \frac{\lambda^{\alpha-2}}{r^\alpha} + \frac{\alpha(\alpha-1)}{2r^2} - \frac{(\alpha-2)}{4r^3}[4\alpha m + \alpha(r-2m)], \tag{14b}$$

$$8\pi\tilde{q}_a = (A^{-1}\frac{\alpha\dot{m}}{r^2})\,\hat{r}_a. \tag{14c}$$

We note that

$$8\pi(\tilde{\rho} + \tilde{p}_r) = -A^{-1}\frac{2\alpha\dot{m}}{r^2} - \frac{\alpha(\alpha-2)}{2r^3}(r-2m).$$

In the static case the string equation of state $\tilde{\rho} + \tilde{p}_r = 0$ holds on the horizon for all α. For mass $m(r)$ which falls off at least as fast as $O(1/r^2)$, the string equation of state also holds at future null infinity.

The metric function $m(u,r)$ measures the system mass only for $\alpha = 2$, the GK metric. In general, spherical symmetry allows the mass within two-surfaces of constant u and r to be invariantly defined by the sectional curvature [9] of those surfaces:

$$\begin{aligned} -2M/r^3 \ :&= R_{abcd}\hat{\vartheta}^a\hat{\varphi}^b\hat{\vartheta}^c\hat{\varphi}^d \\ &= -\frac{\alpha^2 m}{2r^3} + \frac{\alpha^2}{4r^2} - \frac{\lambda^{\alpha-2}}{r^\alpha}. \end{aligned} \tag{15}$$

When $\alpha = 2$ then $M = m$.

III. DIFFUSION AND SCALE

A. Diffusion

Continuous diffusive processes can envelope underlying fractal behavior. In the $\alpha = 2$ Schwarzschild atmosphere, the density ρ and the mass function m diffuse according to Eqs.(2) and (3). By analogy with classical diffusion of density in the Schwarzschild spacetime, we assume that the density enveloping fractal behavior obeys

$$\partial_u \Gamma = r^{-\alpha} \partial_r (D r^{\alpha} \partial_r \Gamma), \tag{16}$$

where the mass density Γ is defined on $u = const$ surfaces, using the sectional curvature mass, by the relation $M = \int \Gamma \sqrt{-g} d^3x$ so that

$$4\pi(\Gamma - \Gamma_0) := \lambda^{\alpha-2} r^{-\alpha} \partial_r M. \tag{17}$$

The relationship between the energy density ρ, seen by observer $\hat{v}^a$, in (12b) and the mass density is then given by

$$\begin{aligned} 4\pi\Gamma = 4\pi\rho &+ (\frac{1}{8r^2})[(\lambda/r)^{\alpha-2}\alpha^2(2m' - 1) - 2\alpha(2m' - \alpha + 1)] \\ &+ (\frac{1}{2r^2})(\lambda/r)^{\alpha-2}[(3-\alpha)(\lambda/r)^{\alpha-2} - 1] - \frac{(\alpha-2)}{16r^3}[8\alpha m - \alpha(r - 2m)]. \end{aligned} \tag{18}$$

In the region where $\alpha = 2$, the two densities Γ and ρ are equal.

Assumption (16) is equivalent to assuming a Fickian mass current of the form

$$\partial_u M = 4\pi D(r)\, r^{\alpha} \partial_r \Gamma, \tag{19}$$

and then $\partial_r M = 4\pi\lambda^{2-\alpha} r^{\alpha}(\Gamma - \Gamma_0)$ yields (16). Furthermore, if the sectional curvature mass M diffuses as

$$\partial_u M = r^{\beta} \partial_r (D r^{-\beta} \partial_r M) \tag{20}$$

then $D(r)$ is determined by $\partial_u M$ in (19) and $\partial_r M$:

$$D(r) = D_0 r^{\beta-\alpha}. \tag{21}$$

Note that if $\alpha = 2$ the M diffusion equation is just Eq.(3) with $\beta = 1 - \delta$. For any α, when $D = D_0$, it follows that $\beta = \alpha$ and that M and Γ diffuse reciprocally.

Equation (20) admits the homogeneous solution

$$M_{\rm hom}(r) = M_0 + M_1(r/\lambda)^{1+\alpha} \tag{22}$$

which can be added to each time-dependent solution. There are two important observations that follow from these results. First, the metric of $M_{\rm hom}$ is not asymptotically flat and the diffusive solution would need to be matched to a solution with appropriate asymptotic behavior. Second, the diffusivity depends on a coordinate and not on the size of any correlated mass grouping. This implies that there may be multiple distance scales in the atmosphere.

B. Dimensions

To better understand fractal scales we briefly discuss fractal dimensions [10]. We cannot use the intuitive notion of the number of coordinates needed to locate a point in some region as the dimension of the region, since Peano constructed a curve (1 coordinate) which fills a square (2 coordinates). Consider a point in set Ω, with Ω a subset of an Euclidean space of integer dimension d_E. The notion of topological dimension involves the idea that neighborhoods around a point have boundaries of smaller dimension. Precisely, the topological dimension $d_T(\Omega)$ of set Ω is the smallest integer such that each point has small neighborhoods with boundaries of lesser dimension. Since $d_T(\emptyset) = -1$ it is clear that $d_T(\Omega) \leq d_E$.

The Hausdorff dimension [11] of self-similar sets (fractals) is given in terms of coverings by open sets and is defined as

$$d_H(\Omega) := \lim_{r\to 0} \frac{-\ln N(r)}{\ln r}$$

where $N(r)$ is the smallest number of open balls needed to cover Ω. The definition of d_H is equivalent to the power law $N(r) \approx const \times r^{-d_H}$. When $d_H = 3$, $N(r)$ corresponds to a 3-dimensional Newtonian specific density.

The Hausdorff dimension of a fractal set is not always an integer. The scaling dimension d_S of fractal sets is more intuitive and, since geometric fractals display regular self-similarity, is equal to the Hausdorff dimension:

$$d_S(\Omega) = \frac{\log\,(\text{number of replicas})}{\log(\text{magnification factor})}.$$

Two famous examples are $d_S(\text{Cantor set}) = \frac{\log(2)}{\log(3)}$ in $d_E = 1$, and $d_S(\text{Sierpinski gasket}) = \frac{\log(3)}{\log(2)}$ in $d_E = 2$. An integer valued example is $d_S(\text{Sierpinski gasket}) = \frac{\log(4)}{\log(2)}$ in $d_E = 3$, since all Sierpinski gaskets obey $d_S(\Omega) = \frac{\log\,(d_E+1)}{\log(2)}$. In general, $d_T(\Omega) \leq d_H(\Omega) \leq d_E$.

C. Scales

Maximum scale size is independent of any particular application. Since $\alpha + 1$ is the dimension of the fractal substrate, bounded by d_E, $\alpha_{\max} = 2$.

Consider a vacuum solution at $\alpha = 2$ and ask how it is produced. Beginning at a small scale, one considers the atmosphere to be made of many regions. Within each region an average energy density is found; in some of the regions it will be negative, in some positive, and in some zero. Then a larger region, containing several small regions is considered and again an average energy density is produced until in the end we have the vacuum atmosphere with all of the fluctuations averaging to zero. This process is just a renormalization group averaging procedure and, just as in renormalization group calculations on a lattice, it is necessary at each averaging step to rescale. In lattice calculations, one rescales to maintain the lattice spacing. In an atmospheric calculation with a horizon, the rescaling is done to always keep the physical observers outside the horizon. There are several ways in which this could be accomplished.

One method is to have an observer at constant r examine an area of his 2-sphere at different scales while maintaining the same coordinate value of r at all scales. The entire 2-sphere at $\alpha = 2$ can be tiled by elements of area δA_2. The sum of all the tiles is the total area of the sphere $4\pi r^2$. At small scales, the observer chooses a region of the sphere

containing many area elements to average over. The result of this averaging must be rescaled at each step, as in lattice renormalization, so that the rescaled area is again δA_2. With this averaging procedure, an observer outside the horizon for $\alpha = 2$ will remain outside of the horizon for all α. The $\alpha = 2$ horizon position will persist at all scales.

To see the effect of the rescaling, consider a typical averaging process. Assume that $\alpha = 2$ is the vacuum solution and let the observer look at an area δA_2 and solid angle $\delta\Omega_2$ with $\delta A_2 = r^2\delta\Omega_2$. This region is the result of averaging over larger regions at the lower scale and then rescaling the area size. The larger region would have an area $\delta A_\alpha = \lambda^{2-\alpha} r^\alpha \delta\Omega_\alpha$. The rescaling would adjust $\delta\Omega_\alpha$ to the size of $\delta\Omega_2$ and thereby rescale the areas. The rescaling formulas are

$$\begin{aligned} (\lambda/r)^{2-\alpha}\delta\Omega_\alpha &\longmapsto \delta\Omega_2 \\ \delta A_\alpha &\longmapsto \delta A_2. \end{aligned} \tag{23}$$

The scaling factor $(\lambda/r)^{2-\alpha}$ is expected to be ≤ 1. For metrics without a horizon, the scale factor can be chosen. When there is a horizon, the scale factor is determined by its value on the horizon. To explore the numerical values of this factor in metrics with a horizon, a specific mass solution is needed.

IV. APPLICATIONS

A. Constant mass solution

A specific mass solution is necessary in order to explicitly treat diffusion and fractal behavior. We have seen above that the sectional curvature mass can be written as

$$2M = \frac{\alpha^2}{4}(2m - r) + \lambda^{\alpha-2} r^{3-\alpha}. \tag{24}$$

To illustrate the rescaling process, we explore the case of constant M. For $\alpha = 2$ there is a Schwarzschild vacuum with an horizon at $r = 2m_0$. Assuming that all scales see the horizon

at $2m_0$ we can determine the scaling factor. From Eq.(24) with $M = m_0$, and $r = 2m = 2m_0$ it follows that

$$(2m_0/\lambda)^{\alpha-2} = 1$$

or $\lambda = 2m_0$.

With $m(r)$ from Eq.(24) the fluid parameters in Eq.(14) are

$$8\pi\tilde{q}_a = 0, \tag{25a}$$

$$8\pi\tilde{\rho} = \frac{6(\alpha-2)}{\alpha}(m_0 r^{-3}), \tag{25b}$$

$$8\pi\tilde{p}_r = -\frac{2(\alpha-2)}{\alpha}[m_0 r^{-3} + \lambda^{-2}(\lambda/r)^{\alpha}], \tag{25c}$$

$$8\pi p_\perp = (\frac{\alpha-2}{\alpha})^2[-2m_0 r^{-3} + \lambda^{-2}(\lambda/r)^{\alpha}]. \tag{25d}$$

For this simple example, $\tilde{\rho}$, resulting from fluctuations, is negative [12] and there is no bounding $\tilde{p}_r = 0$ surface. The radial profile describes an atmosphere asymptotically approaching vacuum. If one examines the $\alpha < 2$ metric, it appears not to be asymptotically flat. However, asymptotic flatness applies only to the observer's $\alpha = 2$ spacetime.

B. Solution with finite boundaries

The $M = m_0$ solution is part of the homogeneous solution, Eq.(22). The complete homogeneous solution, modified to produce vacuum Schwarzschild with a horizon at $r = 2m_0$ is

$$M = m_0 + m_1(2-\alpha)(r/\lambda)^{\alpha+1} \tag{26}$$

with

$$m = \frac{4m_0}{\alpha^2} + \frac{4m_1(2-\alpha)}{\alpha^2}(r/\lambda)^{\alpha+1} + \frac{r}{2} - \frac{2\lambda}{\alpha^2}(r/\lambda)^{3-\alpha}. \tag{27}$$

Evaluating this equation on the horizon at $r = 2m_0$ to find the scaling factors gives the equation, with $x = \lambda/2m_0$,

$$1 + (m_1/m_0)(2-\alpha)x^{-(\alpha+1)} - x^{\alpha-2} = 0. \tag{28}$$

This more complex scaling equation has no solution for $m_1 \geq m_0$, and may describe solutions without an horizon. For $m_1 << m_0$ there are two scaling factors. Some typical values for $m_1/m_0 = 0.01$ are

$$\begin{aligned} \alpha &= 7/4, \quad x \simeq 1,\ 1/7.6 \\ \alpha &= 6/4, \quad x \simeq 1,\ 1/11.9 \\ \alpha &= 5/4, \quad x \simeq 1,\ 1/24.5 \\ \alpha &= 1, \quad\ \ x \simeq 1,\ 1/99 \end{aligned} \tag{29}$$

For the complete homogeneous solution the fluid parameters are

$$8\pi\tilde{q}_a = 0, \tag{30a}$$

$$8\pi\tilde{\rho} = \frac{2(\alpha-2)}{\alpha}[3m_0 r^{-3} + (4-5\alpha)(m_1/\lambda^3)(r/\lambda)^{\alpha-2}], \tag{30b}$$

$$8\pi\tilde{p}_r = -\frac{2(\alpha-2)}{\alpha}[m_0 r^{-3} + \lambda^{-2}(\lambda/r)^{\alpha} - 3\alpha(m_1/\lambda^3)(r/\lambda)^{\alpha-2}], \tag{30c}$$

$$8\pi p_\perp = (\frac{\alpha-2}{\alpha})^2[-2m_0 r^{-3} + \lambda^{-2}(\lambda/r)^{\alpha}] + (\frac{\alpha-2}{\alpha})[(5\alpha-4)(2m_1/r^3)(r/\lambda)^{\alpha+1}]. \tag{30d}$$

For solution (27) there are boundaries for some α. The boundary radii follow from $\tilde{p}_r = 0$ at $\lambda = 2m_0$. With $y = r_b/2m_0$,

$$1 + 2y^{3-\alpha} - 3\alpha(m_1/m_0)y^{\alpha+1} = 0. \tag{31}$$

Again choosing $m_1/m_0 = 0.01$, there are boundaries which get larger as α decreases:

$$\begin{aligned} \alpha &= 7/4, \quad y \simeq 11.5 \\ \alpha &= 6/4, \quad y \simeq 45 \\ \alpha &= 5/4, \quad y \simeq 2800 \\ \alpha &= 1, \quad\ \ \text{no boundary.} \end{aligned} \tag{32}$$

Since $\alpha = 2$ is a vacuum solution, one could attempt construction of a matching vacuum for the lower values of α. However, the lack of a boundary at $\alpha = 1$ implies that the boundaries

mark a change from pressure to tension rather than a boundary to vacuum. This supports a picture of an atmosphere which extends farther as the size of the observed fluctuation decreases, all asymptotically approaching vacuum. We again remark that asymptotic flatness applies only to the observer's $\alpha = 2$ spacetime.

C. THE SIERPINSKI GASKET

These ideas can be applied to diffusive processes whose continuum solutions are expected to envelope fractal behavior at smaller scales. As an example, we will use some existing discussions of enveloped fractal structures in flat spaces to discuss the approximate sizes of α and β where $D(r) = D_0 r^{\beta-\alpha} = D_0 r^{\Theta}$. The numerical parameters of interest are α, the diffusivity factor Θ, the Euclidean embedding dimension d_E in which the fractal substrate is embedded, and a fractal dimension d_F (which may not be d_H). Alexander and Orbach [6] and O'Shaughnessy and Procaccia [5] provide some numerical values of d_F and Θ for various fractal processes. The fractal dimension d_F is defined through the diffusion equation which is written as

$$\frac{\partial \Gamma}{\partial u} = \frac{1}{r^{d_F-1}} \frac{\partial (D_0 r^{-\Theta} r^{d_F-1} \partial_r \Gamma)}{\partial r}$$

In $d_E = 3$, the fractal dimension for the Sierpinski gasket is $d_F = 2$ and $\Theta = 0.59$. The corresponding values of α and β are $\alpha = 1,\ \beta = 1.585$. For $d_E = 4$, the fractal dimension is $d_F = 2.31$ and $\Theta = 0.81$. We find $\alpha = 1.32$ and $\beta = 2.13$.

For all of the above examples, we could find a u-independent solution of the diffusion equations as

$$\Gamma = \Gamma_0 r^{1-\beta},$$

$$M = M_0 + M_1 r^{\alpha+1}.$$

The mass scaling is set entirely by the metric parameter α, while the density depends on the scaling of the diffusivity.

V. CONCLUSION

Introducing a parameter, α, into the extended Schwarzschild metric, produces an atmosphere with processes that can be linked to a fractal mass distribution. Specific values of α can be associated with particular fractal substrates in a small scale region of the atmosphere. The continuous solutions of an anomalous diffusion equation in the atmosphere envelope smaller scale fractal processes. Anomalous diffusion can sample fractal processes on a topological level, giving information about the connectivity as well as the fractal dimension. Fluid parameters from the field equations can be studied for a range of α, allowing the effects of fluctuations from one scale to another to be examined.

Other models are possible. For example, a fractal Schwarzschild cell could be used in a universe model like the Lindquist-Wheeler model [13], perhaps describing the fractal galactic density distribution. If the galactic distribution can be modeled by a single fractal process in which both mass and density diffuse, then on time scales that are short compared with the diffusion time, the mass would scale with the fractal dimension while the scaling of the density would involve topological elements of the fractal substrate. In this paper only a single level of scale reduction has been studied. It is possible that an ensemble of solutions, a multifractal set summed over α, could be used to model more complex physical systems.

Another direction for future work is a possible link between fractal atmospheric structure and chaos. A variety of perturbations to a black hole solution have been shown to lead to chaotic behavior, most often chaotic particle orbits [14]. For example, Bombelli and Calzetta [15] have shown that gravitational perturbations of the Schwarzschild solution can produce chaotic orbits. In general, the effects of chaos range from unusual production of gravitational waves [16] to a possible brightening of the black hole [17]. Orbiting particles with spin may also exhibit chaotic behavior [18], [19]. A relation between a phase space fractal dimension and chaos has been established for multi black hole spacetimes [20] and for black hole chaotic scattering processes [21], [22]. We have considered an atmosphere that has fractal structure in the physical spacetime. At one scale, the solution can describe an ideal black hole, or

with time dependence and space dependence, a black hole with a two fluid atmosphere. The effect of perturbations of this atmosphere and the effects of the physical fractal structure on particle orbits is an interesting area for investigation.

---

[1] Vaidya P C 1953 *Nature* **171** 260

[2] Glass E N and Krisch J P 1999 *Class. Quantum Grav.* **16** 1175

[3] Ghez R and Langlois W E 1986 *Am. J. Phys.* **54** 646

[4] Glass E N and Krisch J P 2000 *Gen. Rel. Grav.* **32** 735

[5] O'Shaughnessy B and Procaccia I 1985 *Phys. Rev. Lett.* **54** 455

[6] Alexander S and Orbach R 1982 *J. Physique-Lettres* **43** 625

[7] Dekeyser R, Maritan A and Stella A L 1994 *Diffusion Processes: Experiment, Theory, Simulations*: Proceedings of the Vth Max Born Symposium, (Springer-Verlag, Berlin)

[8] Nelson E 1985 *Quantum Fluctuations*, (Princeton University Press, Princeton)

[9] Misner C W and Sharp D H 1964 *Phys. Rev.* **B136** 571

[10] Hughes B D 1995 *Random Walks and Random Environments* Vol 1 (Oxford University Press, Oxford)

[11] Hastings H M and Sugihara G 1994 *Fractals - A User's Guide for the Natural Sciences* (Oxford University Press. Oxford)

[12] Ford L H and Roman T A 1997 *Phys. Rev.* **D55** 2082

[13] Lindquist R W and Wheeler J A 1957 *Rev. Mod. Phys.* **29** 432

[14] Rugh S E 1994 *Chaos in the Einstein Equations - Characterization and Importance*, in *Deterministic Chaos in General Relativity*, Eds. Hobill D, Burd A and Coley A (Plenum Press,

New York)

[15] Bombelli L and Calzetta E 1992 *Class. Quantum Grav.* **9** 2573

[16] Cornish N J and Franke N E 1997 *Phys. Rev.* **D56** 1903

[17] Levin J 1999 *Phys. Rev.* **D60** 064015

[18] Suzuki S and Maeda K 1997 Phys. Rev. **D55** 4848

[19] Suzuki S and Maeda K 2000 Phys. Rev. **D61** 024005

[20] Dettmann C P, Frankel N F and Cornish N J 1994 Phys. Rev. **D50** R618

[21] Frolov A V and Larsen A L 1999 *Class. Quantum Grav.* **16** 3717

[22] deMoura A P S and Letelier P S 1999 *Chaos and Fractals in Geodesic Motions Around a Non-Rotating Black Hole with an External Halo,* preprint chao-dyn/9910035